MEIJER POTATO UK


Accord

For early season fresh chips

Accord is an early variety, producing good yields of bold, uniform tubers, suitable for both the fresh and processing market. Tubers of Accord are oval in shape, with white skin and flesh, and shallow eyes. Due to it's versatility, Accord is suitable for boiling, mashing, baked and chip consumption.

	Accord
Yield	Medium
Underwater Weight	381
Dry Matter %	19
Maturity	Early
Dormancy	Short Medium
Bruising index (*1)	18
Second Growth (*2)	7.3
Common scab (*2)	6.7
Late Blight (*2)	3.1
Tuber Shape	Round Oval/ Oval
Percentage 50/+	85
Percentage 40/70	76
Skin Colour	Light Yellow
Flesh Colour	White
Uniformity	Regular
Eye Depth	Shallow
Internal Defects (%)	6
Hollow Heart (%)	0
Fry Colour frites October (8 °C)	8
Fry Colour frites February (8 °C)	7
Fry Colour frites May (8 °C)	5
Fry Colour frites June (4 °C)	4
Potato Cyst Nemotades Ro1 (*2)	9
Potato Cyst Nemotades Ro2/3 (*2)	7
Potato Cyst Nemotades Pa2 (*2)	6
Potato Cyst Nemotades Pa3 (*2)	
Wart Disease wz 1 (D1) (*3)	
Wart Disease wz 2 (G1) (*3)	
Wart Disease wz 6 (O1) (*3)	





(*1): 0= none, 50 = severe
(*2): 9 = resistant, 1 = very susceptible
(*3): 10 =resistant, 3 = very susceptible

Characteristics

- Round oval white fleshed tubers suitable for the early season chip production.
- Benefits from dual PCN *G.Rostochiensis* resistance.
- Good tolerance to spraing.

Building on potential

Agronomy

Soil Nitrogen Supply*	0 an 1	2, 3 and 4	5 and 6
N input kg/ha	160-210	130-160	90-120

*0=low, 6 = high. please refer to RB209 Nutrient Management Guide for details and guidance.

Soil Supply*	0	1	2	3	4
P input kg/ha	250	210	170	100	0
K input kg/ha	360	330	300	150	0
Mg input kg/ha	120	80	40	0	0

*0=low, 4 = high. please refer to RB209 Nutrient Management Guide for details and guidance.

Storage	n/a	n/a


50 kg Tuber Count	400	600	800	1000	1200	1400	1600
Population/ha	26743	32155	37277	42629	48431	54226	58818
Spacing/cm	40.89	34.01	29.34	25.65	22.58	20.17	18.59
Seed Rate t/ha	3.34	2.68	2.33	2.13	2.02	1.94	1.84
Population/ac	10823	13013	15086	19252	19600	21945	23804
Spacing/in	16.1	13.39	11.55	10.1	8.89	7.94	7.32
Seed Rate t/ac	1.35	1.08	0.94	0.86	0.82	0.78	0.74

Values are based on an average soil type and planting date using a 72" 2 row system with irrigation.
We recommend using our Seed Calculator App and adjusting according to your own situation and experience.

Timing	Pre-em	Post-em
Metribuzin sensitivity	Slightly	Slightly

General advice for best results

- Susceptible to Rhizoctonia, so treating seed at planting with appropriate products is recommend for best results.
- Reletively low numbers, good attetention to seed rates needed in order to achive target marketable fraction.
- Early blight programme required as Accord is susceptible to Alternaria.



For any further information contact your local Meijer representative or visit our website at: www.meijerspotato.co.uk

MEIJER POTATO UK

Bayfield Brecks Barns
Garden Lane
Letheringsett
Holt
Norfolk
NR25 7DZ
Tel: 01263 710602

Every effort has been made to ensure that the advice, information, recommendations and statements given in this management guide are correct. Meijer Seed Potato Ltd and associated companies cannot be held responsible for any damage or loss resulting from their use.